TrajectoryViz: Interactive visualization of treatment trajectories

♣ PRESENTER: Maarja Pajusalu

METHODS

TrajectoryViz can visualize trajectories based on **discrete treatment periods** with following data structure:

SUBJECT_ID, STATE, STATE_START_DATE and STATE_END_DATE.

Based on this it produces an interactive R Shiny application that displays interactive sunburst plot of the sequential ordering of the states and patient level visualizations of the state sequences selected from the sunburst plot.

The patient level sequences can be filtered, shown with the gaps in treatment and without, and aligned by different events. All these visualizations are interactive allowing both quantifying the interesting aspects or zooming into particular patterns.

OMOP formatted database TrajectoryViz relies on Cohort2Trajectory package in R github.com/HealthInformaticsUT/Cohort2Trajectory . This package summarizes multiple cohorts defined in ATLAS into linear event sequences, required by TrajectoryViz.

RESULTS

To illustrate the capabilities of the TrajectoryViz package we consider two applications:

- 1. **Treatment of asthma patients** (based on cohorts from Markus et al) and
- 2. Diagnostic procedures on cervical cancer patients before and after the diagnosis.

In both cases we utilize the data from **Estonian Health Insurance Fund** and **Estonian Health Information System.**




github.com/HealthInformaticsUT/TrajectoryViz

Explore the temporal patterns of the sequences, understand the data better, spot problems with analysis setup and generate novel hypotheses with the help of TrajectoryViz R package.

Use Case 1: Cervical Cancer patients

Zoom in from the Sunburst: the patients with procedure/result sequence starting with

PAP-HSIL-Colposcopy





In this particular set of patients, we can see clustering of the sequences - **keeping the gaps** between prescriptions enables looking at the **length of periods between procedures and results**

Did the patient follow suggested pattern for PAP tests before?


Did the patient
follow suggested
pattern for
PAP tests after
Selected
diagnosis?
There was a
subset of the
patients who

There was a subset of the patients who spent much more time taking the drugs

Use Case 2: Asthma patients

Zoom in from the Sunburst: the patients from SABA and LABA&ICS cohorts with treatment sequence

SABA-LABA&ICS



In this particular set of patients, we can see clustering of the sequences - removing the gaps between prescriptions enables looking at the treatment as a continuous period

SABA LABA&ICS ... (LABA&ICS-1: N= 199)

Times before or after the STATE LABA&ICS-1

Most of the

patients spent

only 1-2

months

(prescriptions)

either of the

drugs






Create discrete sequences with R package

Cohort2Trajectory



Create interactive visualisations with R package

TrajectoryViz



Explore and find answers to Your questions

- Maarja Pajusalu, Marek Oja, Sirli Tamm, Markus Haug, Raivo Kolde. Institute of Computer Science, University of Tartu, Estonia
- maarja.pajusalu@ut.ee
- Aniek F. Markus, Peter R. Rijnbeek, Jan A. Kors, Katia Verhamme. TreatmentPatterns: An R package to analyze treatment patterns of a study population of interest. OHDSI 2020 Global Symposium. ohdsi.org/2020-global-symposium-showcase-92









This article has been financed/supported by European Social Fund via "ICT programme" measure.